

Urokinase 국소 주입법으로 치료한 호산구 증다증에 동반된 대량 심부정맥 혈전증 1례

정인현¹ · 최동훈¹ · 김혜진¹ · 경희두¹ · 구본권¹ · 조승연¹ · 이도연²

A Case of Eosinophilia Associated with Massive Deep Vein Thrombosis Treated with Local Urokinase Infusion

In Hyun Jung, MD¹, Donghoon Choi, MD¹, Hye Jin Kim, MD¹, Hee Doo Kyung, MD¹, Bon-Kwon Koo, MD¹, Seung-Yun Cho, MD¹ and Do Yun Lee, MD²

¹Cardiology Division, Yonsei Cardiovascular Center and Cardiovascular Research Institute,

²Radiology, Yonsei University College of Medicine, Seoul, Korea

ABSTRACT

Venous thrombosis is uncommon in young individuals. Hypereosinophilia is a rare cause of thrombosis that usually involves the heart and small vessels like such as retinal vessels. We report a case of massive deep vein thrombosis that developed in a young patient with hypereosinophilia who was successfully treated with continuous local infusion of urokinase. (**Korean Circulation J 2001;31(2):256-261**)

KEY WORDS : Deep vein thrombosis · Eosinophilia · Urokinase.

서 론

가

가¹⁾ 가³⁾⁴⁾ (post - thrombotic syndrome) 가⁵⁾ 가²⁾ okinase 1 ur -

: 2000 12 4
: 2001 2 28
: , 120 - 752

: (02) 361 - 7049 · : (02) 393 - 2041
E - mail : cdhlyj@yumc.yonsei.ac.kr 36 가

증 례

1 X -

4

(Fig. 1).

가 (popliteal vein)

120/70 mmHg, 36 ,

86/ , 18/ .

(Fig. 2).

sheath Multi - sideport(Cook® Bloomington, USA)

21,500/mm³(35.5%,

16.2%, 45.7%(9,800/mm³),

2.3%), 11.4 g/dl, 337,000/mm³

ESR 26 mm/hr . P time(INR) 1.09, aPTT

38.1 , thrombin time 23.4

ELISA

rheumatoid factor, VDRL

HBsAg, Anti - HBc, HBeAg

HBV - DNA 5609.6 pg/ml . ANA, ANCA,

anti - phospholipid Ab IgG IgM

lupus anticoagulant . Protein C 73%, 5

protein S 84%, anti - thrombin 62% factor V

urokinase 480 가

coumadin 7




Fig. 1. Transthoracic echocardiography reveals no deposition of thrombus or eosinophils within the ventricular cavity.




Fig. 2. Venogram on admission shows multiple filling defects from distal inferior vena cava (A) to both popliteal veins (B, C). IVC : inferior vena cava, RF : right femur, LF : left femur.




Fig. 3. Follow-up venogram after thrombolysis and heparin infusion shows a resolution of the previous multiple filling defects within both iliac and femoral veins. There is a filling defect remaining at distal inferior vena cava (white circle) which is suggestive of organized thrombi.

18 (mural thrombi)
(bronchial arteriol)

(Fig. 3).

glucocorticoid Walker⁸⁾

30

B 가

3

11,660/mm³ 가

coumadin 5 mg 가 .¹⁰⁻¹⁴⁾

Elouaer -

Blanc⁷⁾ 가

고 안 가

50 가

anti - thrombin III, protein 37가

C, protein S , antiphospholipid antibody .¹⁹⁾ (1) 가

1,500/mm³ 6 6

factor V Leiden mutation G20210A

prothrombin gene mutation .¹⁾¹⁵⁾ (2) (3)

methionine cobala -

min, folate, pyridoxine

hyperhomo - cysteinemia

가 .¹⁶⁾ 가

3

(major base protein), 1973

(eosinophilic cationic protein), Spitzer Garson

(eosinophil - derived neurotoxin) 23

가

.¹⁷⁾¹⁸⁾ 가

.²⁰⁾

.⁵⁾ Elouaer - Blanc⁷⁾ lupus anticoagulant가

lupus antico -

(main hepatic vein) agulant가 가

antiphospholipid antibody IgM IgG가 lupus anti - coagulant가 90% aPTT가 (cath -
 phospholipid antibody가 anti - eter directed - thrombolysis)
 lupus anticoagulant 가 (venous patency) 요 약
 postphlebitic syndrome 가 1
 24 urokinase
 15 가
 (Low - mo - 중심 단어 : Urokinase.
 lecular - weight - heparin)
 coumadin K 5
 3
 tissue plasminogen activator
 가 2 4
 (regional infusion)

REFERENCES

- 1) Lensing AW, Prandoni P, Prins MH, Buller HR. *Deep-vein thrombosis. Lancet* 1999;353:479-85.
- 2) Kniffin WD, Baron JA, Barrett J, Birkmeyer JD, Anderson FA. *The epidemiology of diagnosed pulmonary thromboembolism and deep vein thrombosis in the elderly. Arch Intern Med* 1994;154:861-6.
- 3) Kearon C, Crowther M, Hirsh J. *Management of patients with hereditary hypercoagulable disorders. Ann Rev Med* 2000;51:169-85.
- 4) Gurgey A. *Clinical manifestations in thrombotic children with factor V Leiden mutation. Pediatr Hematol Oncol* 1999;16:233-7.
- 5) Weller PF, Bubley GJ. *The idiopathic hypereosinophilic syndrome. Blood* 1994;83:2759-79.
- 6) Ferguson GT, Starkebaum G. *Thromboangiitis obliterans associated with idiopathic hypereosinophilia. Arch Intern Med* 1985;145:1726-8.
- 7) Elouaer-Blanc L, Zafrani ES, Farcet JP, Saint-Marc Girardin FM, Mathieu D, Dhumeaux D. *Hepatic vein obstruction in idiopathic hypereosinophilic syndrome. Arch Intern Med* 1985;145:751-3.
- 8) Walker M. *Idiopathic hypereosinophilia associated with hepatic vein thrombosis. Arch Intern Med* 1987;147:2220-1.
- 9) Schulman H, Hertzog L, Zirkin H, Hertzanu Y. *Cerebral sinovenous thrombosis in the idiopathic hypereosinophilic syndrome in childhood. Pediatr Radiol* 1999;29:595-7.
- 10) Kim KT, Lee IJ, Chung KY, Chun SI. *Hypereosinophilic Syndrome on the Extremities Associated with abnormal Peripheral Blood Circulation. Kor J Dermatol* 1995;33:104-8.
- 11) Chung JP, Nam DK, Lee SJ, Lee EK, Han JS, Ko YW. *A Clinical Study on Eosinophilia; With a Report of 5 Cases of Hypereosinophilic Syndrome. Kor J Hematol* 1988;23:127-37.
- 12) Shin KS, Choi YM, Chae SA, Hyung SM. *A Clinical Study*

- of Cause of Eosinophilia. *Chungbuk Med J* 1996;6:105-14.
- 13) Yim SJ, Hong KW, Oh HJ, Khang KK, Han SS, Kim KH. A Case of Hypereosinophilic Syndrome. *Kor J Hematol* 1983;18:227.
 - 14) Lee SB, Choi OS, Kang HS, Choe CW, Kim KS, Kim MS, et al. A Case of Multifocal Cerebral Infarction Associated with Idiopathic Hypereosinophilic Syndrome. *Kor J Med* 1997;52:419-23.
 - 15) Weinmann EE, Salzman EW. Deep-vein thrombosis. *N Engl J Med* 1994;15:1630-41.
 - 16) den Heijer M, Blom HJ, Gerrits WB, Rosendaal FR, Haak HL, Wijermans PW, et al. Is hyperhomocysteinemia a risk factor for recurrent venous thrombosis? *Lancet* 1995;345:882-5.
 - 17) Mukai HY, Ninomiya H, Ohtani K, Nagasawa T, Abe T. Major basic protein binding to thrombomodulin potentially contributes to the thrombosis in patients with eosinophilia. *Br J Haematol* 1995;90:892-9.
 - 18) Babapulle FB. Clonal eosinophilic disorders and the hypereosinophilic syndrome. *Blood reviews* 1997;11:129-45.
 - 19) Chusid MJ, Dale DC, West BC, Wolff SM. The hypereosinophilic syndrome: Analysis of fourteen cases with review of the literature. *Medicine* 1975;54:1-27.
 - 20) Takai K, Sanada M. Hypereosinophilic syndrome evolving to acute lymphoblastic leukemia. *Int J Hematol* 1991;54:231-9.
 - 21) Esmon NL, Smirnov MD, Esmon CT. Thrombogenic mechanisms of antiphospholipid antibodies. *Thromb Haemost* 1997;78:79-82.
 - 22) Ravel R. *Clinical laboratory medicine*. 6th ed. Louis: Mosby Year book;1995. p.102-3.
 - 23) Harada M, Fujisawa Y, Sakisaka S, Kawaguchi T, Taniguchi E, Sakamoto M, et al. High prevalence of anti-cardiolipin antibodies in hepatitis C virus infection: lack of effect on thrombocytopenia and thrombotic complication. *J gastroenterol* 2000;4:272-7.
 - 24) Hull RD, Raskob GE, Pineo GF, Green D, Trowbridge AA, Elliott CG, et al. Subcutaneous low-molecular-weight-heparin with intravenous standard heparin in proximal deep-vein thrombosis. *N Engl J Med* 1992;326:975-82.
 - 25) Comerota AJ. An overview of thrombolytic therapy for venous thromboembolism. In: Comerota AJ. *Thrombolytic therapy*. Orlando: Grune & Stratton;1988. p.65-89.
 - 26) Bates SM, Hirsh J. Treatment of venous thromboembolism. *Thromb Haemost* 1999;82:870-7.